	记录编号 FN: SMP-RD-A0002-R06	版本号 Ver: 2.1
	归档 Filing: filed immediately, permanently kept	生效日期 Ed: 2023.01.12

产品使用说明书 Product Instruction Manual


编号 No.: V204-00

产品名称 Product Name: Media C-01

主货号 Main Art. No.: LB003

编制 Preparation	编制人 Prepared by	编制日期 Date of preparation
研发部 RD		
审核 Review	审核人 Reviewed by	审核日期 Date of Review
研发部 RD		
生产部 PD		
质量部 QD		
质量部 QD		
批准 Approval	批准人 Approved by	批准日期 Date of Approval
质量部 QD		

日期 Date	编制人 Prepared by	版本号 Version No.	主要修订内容 Main revised content
2024.05.09	惠淑怡 Hui shuyi	00	新增 New file

	记录编号 FN: SMP-RD-A0002-R06	版本号 Ver: 2.1
	归档 Filing: filed immediately, permanently kept	生效日期 Ed: 2023.01.12

产品使用说明书 Product Instruction Manual

多宁/DuoNing

动物细胞高性能培养基 High-Performance Culture Medium for Animal-Cells

V204-00

【产品名称 Product name】Media C-01

【货号 Art. No.】LB003

液体包装 liquid packaging

【产品说明 Product description】

Media C-01 是一种无动物来源成分、无蛋白成分、化学成分限定的基础培养基，适合采用中国仓鼠卵巢细胞（CHO）进行治疗性蛋白产品研发和生产过程中的分批培养、补料分批培养和灌流培养。Media C-01 不含有 L-谷氨酰胺。适合 CHOK1、CHOK1SV、CHOS、DG44 等不同细胞株的培养。

Media C-01 is a chemical defined basal medium with no animal origin components, no protein, which is suitable for batch culture, fed-batch culture and perfusion culture in the development and production of therapeutic protein products by Chinese hamster ovary (CHO). Media C -01 does not contain L- glutamine. It is suitable for the culture of different cell lines, such as CHOK1, CHOK1SV, CHOS, DG44.

【细胞培养 Cell culture】

- ① 建议细胞接种密度 Suggested cell inoculation density: $0.3\sim1.0\times10^6$ cells/mL.
- ② 温度 Temperature: 36.5℃
- ③ CO₂: 6~8%

【细胞驯化 Cell domestication】

多数细胞株使用本产品是不需要任何驯化，直接接种到本培养基，传代三次以上即可。对有些细胞株，使用本系列培养基时可能要采用驯化，具体步骤如下：

Most cell lines use this product without any domestication, and can be directly inoculated into this medium and passed for more than three times. For some cell lines, domestication may be used when using this series of medium, and the specific steps are as follows:


① 直接驯化 Direct domestication

大部分细胞株可以直接驯化至 Media C-01 中。

Most cell lines can be directly domesticated to Media C-01.

细胞接种密度 Cell inoculation density: $3.0\sim8.0\times10^5$ cells/mL

至少传代 2~3 代，倍增时间正常稳定，细胞活率 >90%，表示细胞株驯化完成。

	记录编号 FN: SMP-RD-A0002-R06	版本号 Ver: 2.1
	归档 Filing: filed immediately, permanently kept	生效日期 Ed: 2023.01.12

After at least 2~3 generations, the doubling time is normal and stable, and the cell viability is more than 90%, indicating that the cell strain has been domesticated.

② 连续驯化 Continuous domestication

- 细胞株在原培养基培养至指数生长期中期，细胞活率>90%时，接种到 50%: 50 % (Media C-01: 原培养基) 体积比配制的混合培养基中，接种密度在 $3\sim5\times10^5$ cells/mL，在 37°C 和 6% CO₂ 培养。细胞培养 3~4 天达到 1×10^6 cells/mL 以上，传代；

The cell strain was cultured in the original medium to the middle of exponential growth period, and when the cell viability was more than 90%, it was inoculated into the mixed medium with the volume ratio of 50%: 50% (Media C-01: original medium), and the inoculation density was $3\sim5\times10^5$ cells/mL, and it was cultured at 37°C and 6% CO₂. The cells were cultured for 3~4 days to reach more than 1×10^6 cells/mL, and then subcultured.

- 将细胞接种到 75%: 25% (Media C-01: 原培养基) 体积比配制的混合培养基中，接种密度在 $3\sim5\times10^5$ cells/mL，在 37°C 和 6% CO₂ 培养。细胞培养 3~4 天达到 1×10^6 cells/mL 以上，传代；

Cells were inoculated into a mixed medium with the volume ratio of 75%: 25% (Media C-01: original medium), and the inoculation density was $3\sim5\times10^5$ cells/mL, and cultured at 37°C and 6% CO₂. The cells were cultured for 3~4 days to reach more than 1×10^6 cells/mL, and then subcultured.


- 将该细胞接种到 100% Media C-01 中，接种密度在 $3\sim5\times10^5$ cells/mL，在 37°C 和 6%CO₂ 培养。细胞培养 3~4 天达到 1×10^6 cells/mL 以上，传代；

The cells were inoculated into 100% Media C-01 with the inoculation density of $3\sim5\times10^5$ cells/mL, and cultured at 37°C and 6%CO₂. The cells were cultured for 3~4 days to reach more than 1×10^6 cells/mL, and then subcultured.

- 在 100% Media C-01 中，至少传代 2~3 代，倍增时间正常稳定，细胞活率>90%，表示细胞株驯化完成；
In 100% Media C-01, at least 2~3 generations, the doubling time is normal and stable, and the cell viability is more than 90%, indicating that the cell strain has been domesticated.

- 采用本驯化程序时，若细胞还是生长很慢或活度很低，可考虑从 10: 90 (Media C-01: 原培养基) 体积比配制的混合培养基起，缓慢增加 Media C-01 的比例到 25: 75, 50: 50, 75: 25, 100: 0；或者过程中离心收集细胞，重新进行传代。

When adopting this domestication procedure, if the cells still grow slowly or have low activity, we can consider slowly increasing the ratio of Media C-01 to 25: 75, 50: 50, 75: 25, 100: 0 from the mixed medium prepared with

	记录编号 FN: SMP-RD-A0002-R06	版本号 Ver: 2.1
	归档 Filing: filed immediately, permanently kept	生效日期 Ed: 2023.01.12

a volume ratio of 10: 90 (Media C-01; original medium). Or the cells are collected by centrifugation during the process and subcultured again.

【细胞冻存 Cell cryopreservation】

- ① 在超净工作台上准备冻存液：90% Media C-01 + 10% 二甲基亚砷（DMSO）混合液，2~8℃预冷（DMSO 稀释时会释放热量）；

Prepare frozen solution on the super clean workbench : 90% Media C-01 +10% dimethyl sulfoxide (DMSO) mixed solution, precooling at 2~8℃ (heat will be released when DMSO is diluted);

- ② 冻存细胞液：处于对数生长期，密度大于 1.5×10^6 cells/mL，活率大于 95%，一般建议冻存密度为 $1.0 \sim 1.5 \times 10^7$ cells/mL；

Frozen cell fluid: in the exponential growth period, the density is greater than 1.5×10^6 cells/mL, and the viability is greater than 95%. Generally, it is recommended that the frozen storage density is $1.0 \sim 1.5 \times 10^7$ cells/ml;

- ③ 细胞液 800rpm 离心 5 min；

Cell fluid was centrifuged at 800rpm for 5 min;


- ④ 缓慢倒出上清液，使用冻存液重新悬浮细胞，冻存密度 $1.0 \sim 1.5 \times 10^7$ cells/mL，将细胞转移至无菌冻存管中；

Slowly pour out the supernatant, resuspend the cells with cryopreservation solution, the cryopreservation density is $1.0 \sim 1.5 \times 10^7$ cells/mL, and transfer the cells to a sterile cryopreservation tube;

- ⑤ 将冻存管置于含异丙醇的冻存盒中，-80℃冻存过夜，再转移至液氮罐中长期贮存。如果没有冻存盒，可手动梯度降温，步骤如下：

Place the cryopreservation tube in the cryopreservation box containing isopropyl alcohol, freeze it at - 80 °C overnight, and then transfer it to the liquid nitrogen tank for long-term storage. If there is no freezing box, the temperature can be reduced manually by gradient as follows:

- 4℃冻存 30min；
- freeze at 4℃ for 30min；
- -20℃冻存 2~4 小时；
- freeze at -20℃ for 2~4h；
- -80℃冻存过夜；
- freeze at - 80℃ overnight；

	记录编号 FN: SMP-RD-A0002-R06	版本号 Ver: 2.1
	归档 Filing: filed immediately, permanently kept	生效日期 Ed: 2023.01.12

- 转移至液氮罐中长期贮存。
- transfer frozen cells to liquid nitrogen tank for long-term storage.

【细胞复苏 Cell resuscitation】

- ① 准备 36.5℃ 温水，用于解冻细胞；

Prepare a 36.5 °C warm water to thaw frozen cells;

- ② 准备 15 ml 无菌离心管，加入 2~5mL 的 Media C-01；

Prepare 15 ml sterile centrifuge tube and add 2~5mL Media C-01;

- ③ 从液氮罐中取出冻存管，迅速在 36.5℃ 水浴锅中将细胞融化；

Take out the frozen tube from the liquid nitrogen tank and quickly thaw frozen cells in 36.5°C warm water;

- ④ 用 75% 的乙醇擦拭冻存管后，在无菌操作台中打开冻存管，将细胞液转移至含 2~5 mL 的 Media C-01 的 15 ml 离心管中，吹打混匀，800rpm 离心 5 min；

After wiping the cryopreservation tube with 75% ethanol, open the cryopreservation tube in the sterile operation table, transfer the cell fluid to a 15 ml centrifuge tube containing 2-5 mL of Media C-01, blow and mix well, centrifuge at 800 rpm for 5 minutes;

- ⑤ 缓慢倒出上清液，使用 20~30 ml 预热 Media C-01 重新悬浮，转移至 125 ml 摇瓶中；

Slowly pour out the supernatant, resuspend with 20~30 ml preheated Media C-01, and transfer to a 125 ml shake flask;

- ⑥ 放置于 36.5℃，6~8% CO₂，80% 湿度，110~130rpm 的摇床中培养；

Place it in a shaking incubator with 6~8% CO₂, 110 ~ 130rpm, at 36.5°C for culture;


- ⑦ 培养 2~3 天后，对细胞进行计数传代。

After 2~3 days of culture, the cells were counted and subcultured.

【细胞传代 Cell passage】

按照 5E5 ~6E5 的密度进行传代，每隔 2~3 天计数，传代。前 3 次传代，体积不变，以恢复细胞活力。待细胞活力恢复正常，达 90% 以上后，以 5E5 ~6E5 的密度进行扩增，直至达到所需种子体积，种子状态正常的标准：活力大于 95%，细胞形态规则圆整，生长倍增时间正常。

The cells are seeded at 5E5 ~6E5, count and subculture every 2 ~ 3 days. In the first three passages, the volume remained unchanged to restore cell viability. After the cell viability recovers to normal and reaches more than 90%. The seed cells were expanded at the density of 5E5 ~6E5 until reaching the required volume. The criteria for normal seed state: the viability was greater than 95%, the cell morphology was regular and round, and the growth doubling time was normal.

	记录编号 FN: SMP-RD-A0002-R06	版本号 Ver: 2.1
	归档 Filing: filed immediately, permanently kept	生效日期 Ed: 2023.01.12

【储存和有效期 Storage and validity period】

Media C-01 液体包装: 2~8℃避光储存, 有效期为 12 个月。

Media C-01 liquid packaging: 2℃ to 8℃, protect from light; Shelf life: 12 months.

【生产企业信息 Manufacturer information】

名称 Name: 无锡多宁生物科技有限公司 Wuxi Duoning Biotechnology co.,Ltd

地址 Address: 无锡新加坡工业园新集路 2-1、2-2 号厂房 No.2-1, No.2-2, Xinji Road, Singapore Industrial Park, Wuxi

电话 Tel: 0510-85956600

网址 Website: www.duoningbio.com