	记录编号 FN: SMP-RD-A0002-R06	版本号 Ver: 2.1
	归档 Filing: filed immediately, permanently kept	生效日期 Ed: 2023.01.12

产品使用说明书 Product Instruction Manual

多宁/DuoNing

动物细胞高性能培养基 High-Performance Culture Medium for Animal-Cells

V111-01

【产品名称 Product name】高效补料培养基 B High-efficiency feed medium B

【货号 Art. No.】BP100-001、BP100-005、BP100-015、BP100-030、BP100-1kg-01、BP100-100g-02、BP100-1kg-02、BP100-5kg-02、BP100-10kg-02、BP100-25kg-02 粉末包装 Powder packaging

【产品说明 Product description】


高效补料培养基 B 是搭配 2×103A 高效补料培养基 A、2×104 高效补料培养基 A、2×105 高效补料培养基 A、2×107 高效补料培养基 A、103 高效补料培养基 A 使用。其中，2×103A 高效补料培养基是一种无动物来源成分、无蛋白的高效补料培养基，适用于采用中国仓鼠卵巢细胞（CHO）进行治疗性蛋白产品研发和生产过程中的补料分批（fed-batch）培养。本产品成分不含有次黄嘌呤、胸腺嘧啶、L-谷氨酰胺，不含有 P188 和碳酸氢钠。适合采用 GS 系统和 DHFR 筛选系统的 CHO-K1、CHO-S、DG44、CHOZN 等不同细胞株的培养。2×103A 高效补料培养基分为 2×103A 高效补料培养基 A 和高效补料培养基 B 两个组分。

The high-efficiency feed medium B is used together with 2×103A high-efficiency feed medium A, 2×104 high-efficiency feed medium A, 2×105 high-efficiency feed medium A, 2×107 high-efficiency feed medium A and 103 high-efficiency feed medium A. Among them, 2×103A high-efficiency feed medium is a kind of high-efficiency fed-batch culture medium without animal-derived components and protein, which is suitable for the research and development of therapeutic protein products by Chinese hamster ovary (CHO) and the fed-batch culture in the production process. This product contains no hypoxanthine, thymine, L- glutamine, P188 and sodium bicarbonate. It is suitable for the culture of different cell lines, such as CHO-K1, CHO-S, DG44 and CHOZN, using GS system and DHFR screening system. 2×103A high-efficiency feed medium is divided into 2×103A high-efficiency feed medium A and high-efficiency feed medium B.

【使用指南 User guide】

2×103A 高效补料培养基是为提高 CHO 细胞生长和生产性能专门设计的一种高效补料配方，其含有丰富的氨基酸和维生素等营养物质，可以满足细胞高密度培养和产物高表达时的营养需求。结合细胞株特性，采用合理的补料策略可以明显改善细胞的生长和抗体的表达量。

2×103A high-efficiency feed medium is an efficient feeding formula specially designed to improve the growth and production performance of CHO cells. It is rich in nutrients such as amino acids and vitamins, which can meet the nutritional needs of high-density culture of cells and high expression of products. Combined with the characteristics of cell

	记录编号 FN: SMP-RD-A0002-R06	版本号 Ver: 2.1
	归档 Filing: filed immediately, permanently kept	生效日期 Ed: 2023.01.12

lines, reasonable feeding strategy can obviously improve the growth of cells and the antibody expression.

推荐的补料策略:

Recommended feeding strategy:

1. 第一次补料建议在细胞指数生长期中后期, 通常细胞密度达至  $2.0 \times 10^6$  cells/mL 以上, 开始补料。建议补料量为: 第 3 天建议 5%、第 5 天建议 5%、第 7 天建议 7.5%、第 9 天建议 7.5%、第 11 天建议 5%、第 13 天建议 5%。

It is suggested that the first feeding should be started in the middle and late period of cell exponential growth, when the cell density is above  $2.0 \times 10^6$  cells/mL. The recommended feeding amount is: 5% on the 3rd day, 5% on the 5th day, 7.5% on the 7th day, 7.5% on the 9th day, 5% on the 11th day and 5% on the 13th day.

2. 建议总补液量不超过 40%。

It is suggested that the total feeding amount should not exceed 40%.

3.  $2 \times 103A$  高效补料培养基可以与 BR7 组合使用, 培养效能可进一步提高, BR7 每次添加浓度推荐为 1%。

$2 \times 103A$  high-efficiency feed medium can be used in combination with BR7, and the culture efficiency can be further improved. The recommended concentration of BR7 for each addition is 1%.

注意事项 Precautions:

① 培养 GS 筛选系统工程细胞株时, 根据需要添加次黄嘌呤和胸腺嘧啶。

Hypoxanthine and thymine were added as needed when cultivating GS screening system engineering cell lines.

② 培养 DHFR 筛选系统工程细胞株时, 根据需要添加 3~8mM L-谷氨酰胺。

3~8mM L- glutamine was added as needed when cultivating DHFR screening system engineering cell lines.

③ 本产品液体包装在 2~8℃ 放置过程会产生部分沉淀, 因此建议两个月内使用完。

The liquid packaging of this product will partially precipitate when it is placed at 2~8℃, so it is recommended to use it within two months.

## 【配制指南 Preparation guide】

适用于粉末包装 (以 1L 为例)


Suitable for powder packaging (taking 1L as an example)

1. 准备配液体积 80% 的超纯水 (35~40℃);

Prepare ultrapure water (35 ~ 40℃) with 80% of the liquid volume;

2. 加入  $2 \times 103A$  高效补料培养基 A 粉末 99.03g, 搅拌 20 min, 溶解完全, 此时溶液略有浑浊, pH 在 4.0 左右;

Add 99.03g of  $2 \times 103A$  high-efficiency feed medium A powder, stir for 20 min, and dissolve completely. At this time, the solution is slightly turbid, and the pH is about 4.0;

	记录编号 FN: SMP-RD-A0002-R06	版本号 Ver: 2.1
	归档 Filing: filed immediately, permanently kept	生效日期 Ed: 2023.01.12

3. 另外准备 50ml 0.4M NaOH 溶液，加入高效补料培养基 B 粉末 2.96g，搅拌 10 min，溶解完全，记为 B 溶液；  
In addition, prepare 50 ml of 0.4M NaOH solution, add 2.96g of high-efficiency feed medium B powder, stir for 10 min, and record it as solution B;

4. 加入 B 溶液，继续搅拌 20 min；  
Add solution B and continue stirring for 20 min;

5. 调节 pH 至 6.70 左右，溶液澄清；  
Adjust the pH to about 6.70, and the solution will be clarified;

6. 加入 2.1g 碳酸氢钠，调节 pH 至 6.90~7.10；  
Add 2.1g sodium bicarbonate to adjust the pH to 6.90 ~ 7.10;

7. 定容，搅拌 5~10 min；  
Constant volume, stirring for 5 ~ 10 min;

8. 用 0.22μm 过滤器除菌过滤；  
Sterilize and filter with a 0.22μm filter;

9. 2×103A 培养基液体渗透压范围在 750~850 mOsm/kg。  
The osmotic pressure of 2×103A medium is 750~850 mOsm/kg.

#### 【储存、有效期或复验期 Storage condition, validity period or retest date】

上海生产基地，干粉包装：2~8℃避光储存，有效期为 24 个月。

Shanghai production base, powder packaging: 2°C to 8°C, protect from light; validity period: 24 months.

无锡生产基地，干粉包装：2~8℃避光储存，复验期为 24 个月。

Wuxi production base, powder packaging: 2°C to 8°C, protect from light; retest date: 24 months.

#### 【生产企业信息 Manufacturer information】

名称 Name: 上海多宁生物科技股份有限公司 Shanghai Duoning Biotechnology Co., Ltd.

地址 Address: 上海市松江区民强路 1525 弄 30 幢 Building 30, Lane 1525, Minqiang Road, Songjiang District, Shanghai

电话 Tel: 021-64340155

网址 Website: [www.duoningbio.com](http://www.duoningbio.com)

名称 Name: 无锡多宁生物科技有限公司 Wuxi Duoning Biotechnology co.,Ltd

地址 Address: 无锡新加坡工业园新集路 2-1、2-2 号厂房 No.2-1, No.2-2, Xinji Road, Singapore Industrial Park, Wuxi

电话 Tel: 0510-85956600

网址 Website: [www.duoningbio.com](http://www.duoningbio.com)